Masimo LiDCO[®] Hemodynamic Monitoring System



The LiDCO Haemodynamic Monitoring System provides beat-to-beat advanced haemodynamic monitoring to support informed decision making in high-acuity care areas such as the Operating Room.

- > Uses existing arterial line and blood pressure transducer to monitor haemodynamic parameters
- > PulseCO[™] algorithm converts beat-to-beat blood pressure into its constituent parts, flow and resistance, scaled to each patient's age, height, and weight
- > Proven to be reliable on patients on vasoactive drugs¹



Key Features

Trend Notifications

Internal Battery

For portability around the bed space and seamless transition to different clinical areas

Day/Night Mode

Switch between day and night mode to best suit your environment

Guided Protocols

To help you assess fluid responsiveness (Fluid Challenge, Passive Leg Raise and New Ventilator Tests)

Education

On-screen educational screens for calibration

Preload Response

Displays volume status indications for Pulse Pressure Variation (PPV%) and Stroke Volume Variation (SVV%)

Alerts user to significant haemodynamic changes (>10%) to encourage an immediate response to patient deterioration

Short-term Trend

2-minute window for greater focus on the immediate response to interventions

Long-term Trend

Facilitates interpretation of trends from the start of monitoring, which can be customised to show only the parameters you need

Event Response

Allows you to mark and monitor specific events, like a fluid challenge



Easy Setup and Operation

The LiDCO Monitor is designed for efficient setup and simple operation, with an intuitive, easy-to-interpret display—-facilitating effective haemodynamic management even on those patients who are haemodynamically unstable and require fluid and drug support.




- > Plug-and-play operations using the invasive blood pressure output port on the vital signs monitor
- > Monitor using the existing blood pressure transducer, eliminating the need for an additional disposable

Reductions in Postoperative Complications and Costs

In a randomised, controlled trial of 743 patients undergoing major abdominal surgery, researchers found haemodynamic optimisation with LiDCO led to a 20% reduction in postoperative complications and, as a result, patients monitored with LiDCO were on average \$530 less expensive to treat than control patients who were not monitored.²

Reductions in 30-Day and 180-Day Mortality

In a study comparing the outcomes of 600 emergency laparotomy patients, researchers found that, following the implementation of a program including LiDCO technology, there was a significant decrease in mortality at 30 days (from 21.8 to 15.5%) and 180 days (from 29.5 to 22.2%).³



Parameters and Indicators

The beat-to-beat parameters displayed by the LiDCO monitor provide immediate feedback on a patient's fluid and haemodynamic status.

The LiDCO monitor provides the following parameters:

- > Stroke Volume (SV): The amount of blood pumped by the left ventricle of the heart in one contraction
- > Cardiac Output (CO): The amount of blood the heart pumps through the circulatory system in a minute, calculated by multiplying the stroke volume by the patient's heart rate
- > Systemic Vascular Resistance (SVR): Reflects the resistance to flow and is calculated as the quotient of pressure and cardiac output
- > Oxygen Delivery (DO2): The amount of oxygen delivered to the tissues, calculated as the product of cardiac output and oxygen concentration
- > Stroke Volume Variation (SVV): A dynamic variable that can predict fluid responsiveness in mechanically ventilated patients, SVV is the variation in stroke volume across at least one respiratory cycle
- > Pulse Pressure Variation (PPV): Another dynamic variable that can predict fluid responsiveness in mechanically ventilated patients, PPV is the variation in arterial pulse pressure across at least one respiratory cycle

Monitor Specifications

PHYSICAL CHARACTERISTICS	ORDERING INFORMATION
Weight	LiDCO Hemodynamic Monitor Kit PN 99026
ENVIRONMENTAL	PARAMETERS SUPPORTED
Operating Temperature	Stroke Volume (SV) Cardiac Output (CO) Systemic Vascular Resistance (SVR) Oxygen Delivery (DO2) Stroke Volume Variation (SVV)

Pulse Pressure Variation (PPV)

LiDCO is not licensed for sale in Canada.

¹ LiDCO data on file. ² Pearse R et al. Effect of a perioperative, cardiac output-guided hemodynamic therapy algorithm on outcomes following major gastrointestinal surgery: a randomized clinical trial and systematic review. *JAMA* 2014; 311(21):2181-90. ³ Tengberg LT et al. Multidisciplinary perioperative protocol in patients undergoing acute high-risk abdominal surgery. *Br J Surg* 2017; 104:463-471.

For professional use. See instructions for use for full prescribing information, including indications, contraindications, warnings, and precautions.

Masimo U.S. tel 1-877-4-Masimo info-america@masimo.com Masimo International tel +41-32-720-1111 info-international@masimo.com

